

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
3 July 2003 (03.07.2003)

PCT

(10) International Publication Number
WO 03/053945 A2

(51) International Patent Classification⁷: **C07D 265/36**,
A61K 31/395, A61P 43/00, C07D 209/14, 215/12, 413/12

(74) Agent: **GIDDINGS, Peter, John**; GlaxoSmithKline,
Corporate Intellectual Property CN925.1, 980 Great West
Road, Brentford, Middlesex TW8 9GS (GB).

(21) International Application Number: **PCT/GB02/05812**

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC,
VN, YU, ZA, ZM, ZW.

(22) International Filing Date:
19 December 2002 (19.12.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0130550.7 20 December 2001 (20.12.2001) GB

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK,
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*):
SMITHKLINE BEECHAM P.L.C. [GB/GB]; 980
Great West Road, Brentford, Middlesex TW8 9GS (GB).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **RAMI, Harshad**,
Kantilal [GB/GB]; GlaxoSmithKline, New Frontiers Sci-
ence Park South, Third Avenue, Harlow, Essex CM19 5AW
(GB). **THOMPSON, Mervyn** [GB/GB]; GlaxoSmithK-
line, New Frontiers Science Park South, Third Avenue,
Harlow, Essex CM19 5AW (GB).

Published:


— without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.



WO 03/053945 A2

(54) Title: NOVEL COMPOUNDS



(I)

(57) Abstract: Compounds of formula (I) or a pharmaceutically acceptable salt or solvate thereof: (I) wherein, R¹, R², P, p and n are as defined in the specification, a process for preparing such compounds, a pharmaceutical composition containing such compounds and the use of such compounds in medicine.

Novel Compounds

This invention relates to novel compounds, especially urea derivatives,
5 having pharmacological activity, processes for their preparation, to
compositions containing them and to their use in the treatment of various
disorders.

Vanilloids are a class of natural and synthetic compounds which are
10 characterised by the presence of a vanillyl (3-hydroxy 4-methoxyphenyl) group
or a functionally equivalent group. Vanilloid Receptor (VR1), whose function is
modulated by such compounds, has been widely studied and is extensively
reviewed by Szallasi and Blumberg (The American Society for Pharmacology
and Experimental Therapeutics, 1999, Vol. 51, No. 2.).

15

A wide variety of Vanilloid compounds of different structures are
known in the art, for example those disclosed in European Patent Application
Numbers EP 0 347 000 and EP 0 401 903, UK Patent Application Number
GB 2226313 and International Patent Application, Publication Number WO
20 92/09285. Particularly notable examples of vanilloid compounds or vanilloid
receptor modulators are capsaicin, namely trans 8-methyl-N-vanillyl-6-
nonenamide, isolated from the pepper plant, capsazepine (Tetrahedron, Vol.
53, No. 13, pp. 4791- 4814, 1997) and olvanil - N-(3-methoxy-4-hydroxy-
benzyl)oleamide (J. Med. Chem. 1993, 36, 2595-2604).

25

US Patent Numbers US 3,424,760 and US 3,424,761 both describe a
series of 3-Ureidopyrrolidines that are said to exhibit analgesic, central
nervous system, and psychopharmacologic activities. These patents
specifically disclose the compounds 1-(1-phenyl-3-pyrrolidinyl)-3-phenyl urea
30 and 1-(1-phenyl-3-pyrrolidinyl)-3-(4-methoxyphenyl)urea respectively.


International Patent Applications, Publication Numbers WO 02/08221,
WO 02/16317, WO 02/16318 and WO 02/16319 each disclose certain

vanilloid receptor antagonists and their use in the treatment of diseases associated with the activity of the vanilloid receptor.

Co-pending International Patent Application Number PCT/EP02/04802

5 discloses a series of urea derivatives and their use in the treatment of diseases associated with the activity of the vanilloid receptor.

A structurally novel class of compounds has now been found which also possess Vanilloid receptor (VR1) antagonist activity. The present
10 invention therefore provides, in a first aspect, a compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof:



15 (I)


wherein,

P is phenyl, naphthyl or heterocyclyl;

20 R¹ is selected from -H, halo, alkyl, alkoxy, cycloalkyl, aralkyl, aralkoxy, cycloalkylalkyl, cycloalkylalkoxy, -CN, -NO₂, -OH, -OCF₃, -CF₃, -NR⁵R⁶, -S(O)ₘR⁷, -S(O)₂NR⁵R⁶, -OS(O)₂R⁷, -OS(O)₂CF₃, -O(CH₂)ₓNR⁵R⁶, -C(O)CF₃, -C(O)alkyl, -C(O)cycloalkyl, -C(O)aralkyl, -C(O)Ar, -C(O)(CH₂)ₓOR⁷, -C(O)(CH₂)ₓNR⁵R⁶, -C(O)alkoxy, -C(O)NR⁵R⁶, -
25 (CH₂)ₓC(O)alkoxy, -(CH₂)ₓOC(O)R⁷, -(CH₂)ₓOR⁷, -(CH₂)ₓR⁵R⁶, -(CH₂)ₓC(O)NR⁵R⁶, -(CH₂)ₓN(R⁵)C(O)R⁷, -(CH₂)ₓS(O)₂NR⁵R⁶, -(CH₂)ₓN(R⁵)S(O)₂R⁷, -ZAr, -(CH₂)ₓS(O)₂R⁷, -(OCH₂)ₓS(O)₂R⁷, -N(R⁵)S(O)₂R⁷, -N(R⁵)C(O)R⁷, -(CH₂)ₓN(R⁵)S(O)₂R⁷, -(CH₂)ₓN(R⁵)C(O)R⁷ or -(CH₂)ₓC(O)alkyl;

30

R² is the group:



X is a bond, C, O or NR⁸;

5

R³ is selected from -H, halo, alkyl, alkoxy, cycloalkyl, aryl, aralkyl, aralkoxy, cycloalkylalkyl, cycloalkylalkoxy, -CN, -NO₂, -OH, -OCF₃, -CF₃, -NR⁵R⁶, -S(O)_mR⁷, -S(O)₂NR⁵R⁶, -OS(O)₂R⁷, -OS(O)₂CF₃, -O(CH₂)_xNR⁵R⁶, -C(O)CF₃, -C(O)alkyl, -C(O)cycloalkyl, -C(O)aralkyl, -C(O)Ar, -

10 C(O)(CH₂)_xOR⁷, -C(O)(CH₂)_xNR⁵R⁶, -C(O)alkoxy, -C(O)NR⁵R⁶, -(CH₂)_xC(O)alkoxy, -(CH₂)_xOC(O)R⁷, -(CH₂)_xOR⁷, -(CH₂)_xR⁵R⁶, -(CH₂)_xC(O)NR⁵R⁶, -(CH₂)_xN(R⁵)C(O)R⁷, -(CH₂)_xS(O)₂NR⁵R⁶, -(CH₂)_xN(R⁵)S(O)₂R⁷, -ZAr, -(CH₂)_xS(O)₂R⁷, -(OCH₂)_xS(O)₂R⁷, -N(R⁵)S(O)₂R⁷, -N(R⁵)C(O)R⁷, -(CH₂)_xN(R⁵)S(O)₂R⁷, -(CH₂)_xN(R⁵)C(O)R⁷
 15 or -(CH₂)_xC(O)alkyl;

R⁴ is hydrogen or alkyl;

R⁵ and R⁶ may be the same or different and represent H or alkyl or R⁵ and

20 R⁶ together with the atoms to which they are attached form a C₃₋₆ azacycloalkane, C₃₋₆(2-oxo)azacycloalkane ring or C₅₋₈ polymethylene chain optionally interrupted by heteroatoms such as O or -NR⁸;

Z is O, S or NR⁸;

25

R⁷ is alkyl or aryl;

R⁸ is hydrogen, alkyl or aryl;

n is 2, 3, 4, 5 or 6;

p is 0, 1, 2, 3 or 4;

5 q is 0, 1, 2 or 3;

r is 0, 1 or 2; and


x is 0, 1, 2, 3, 4, 5 or 6.

10

Suitably, P is phenyl, naphthyl, cinnoliny or isoquinoliny. When P is naphthyl, a preferred group is naphth-1-yl. Preferably, P is phenyl.

Suitably, R¹ is halo, alkyl, alkoxy, -C(O)alkyl, -NO₂, -CF₃, -CN or -
 15 OCF₃. More suitably, R¹ is halo, alkyl, -C(O)alkyl or -OCF₃. Preferably, R¹ is halo, -C(O)Me or -OCF₃.

Suitably, R² is



20

Preferably, R² is dihydroindolyl, tetrahydroquinoliny or dihydrobenzo[1,4]oxazinyl. Most preferably, R² is 2,3-dihydroindol-1-yl, 3,4-dihydro-2H-quinolin-1-yl or 2,3-dihydrobenzo[1,4]oxazin-4-yl.

25

Suitably, R³ is halo, alkyl, alkoxy, -CF₃, -CN or aryl. More suitably, R³ is halo or alkyl. Preferably, R³ is fluoro or methyl. Most preferably, R³ is a methyl or fluoro substituted at either the 4- or 5-position on the dihydroindole ring, a methyl group substituted on the 6-position of the dihydroquinoliny ring or a methyl group substituted on the 7-position of the

30

dihydrobenzo[1,4]oxazinyl ring.

Suitably, R⁴ is alkyl. Preferably, R⁴ is methyl.

Suitably, R⁵ is alkyl. Preferably, R⁵ is methyl.

5

When p is 2 or 3 the groups R¹ may be the same or different.
Preferably, p is 1 or 2.

When r is 2 the groups R⁴ may be the same or different.
10 Preferably, r is 0 or 1.

Preferably, n is 2 or 3. Most preferably, n is 2.

Preferably, q is 1 or 2. Most preferably, q is 1.
15

Preferably, x is O.

Particularly preferred compounds according to the invention include
examples E1 to E58 or a pharmaceutically acceptable salt or solvate thereof.
20

The compounds of the formula (I) can form acid addition salts with
acids, such as conventional pharmaceutically acceptable acids, for example
maleic, hydrochloric, hydrobromic, phosphoric, acetic, fumaric, salicylic, citric,
lactic, mandelic, tartaric and methanesulphonic.

25

Compounds of formula (I) may also form solvates such as hydrates,
and the invention also extends to these forms. When referred to herein, it is
understood that the term 'compound of formula (I)' also includes these forms.

30 Certain compounds of formula (I) are capable of existing in
stereoisomeric forms including diastereomers and enantiomers and the
invention extends to each of these stereoisomeric forms and to mixtures
thereof including racemates. The different stereoisomeric forms may be

separated one from the other by the usual methods, or any given isomer may be obtained by stereospecific or asymmetric synthesis. The invention also extends to any tautomeric forms and mixtures thereof.

5 As indicated above, the compounds of formula (I) can form salts, especially pharmaceutically acceptable salts. Suitable pharmaceutically acceptable salts are those use conventionally in the art and include those described in *J. Pharm. Sci.*, 1977, **66**, 1-19, such as acid addition salts.

10 Suitable pharmaceutically acceptable salts include acid addition salts.

 Suitable pharmaceutically acceptable acid addition salts include salts with inorganic acids such, for example, as hydrochloric acid, hydrobromic acid, orthophosphoric acid or sulphuric acid, or with organic acids such, for
15 example as methanesulphonic acid, toluenesulphonic acid, acetic acid, propionic acid, lactic acid, citric acid, fumaric acid, malic acid, succinic acid, salicylic acid, maleic acid, glycerophosphoric acid or acetylsalicylic acid.

 The salts and/or solvates of the compounds of the formula (I) which
20 are not pharmaceutically acceptable may be useful as intermediates in the preparation of pharmaceutically acceptable salts and/or solvates of compounds of formula (I) or the compounds of the formula (I) themselves, and as such form another aspect of the present invention.

25 The compounds of formula (I) may be prepared in crystalline or non-crystalline form, and if crystalline, may be optionally hydrated or solvated. This invention includes in its scope stoichiometric hydrates as well as compounds containing variable amounts of water.

30 Suitable solvates include pharmaceutically acceptable solvates, such as hydrates.

Solvates include stoichiometric solvates and non-stoichiometric solvates.

As used herein the term "alkyl" as a group or part of a group refers to a
5 straight or branched chain saturated aliphatic hydrocarbon radical containing
1 to 12 carbon atoms, suitably 1 to 6 carbon atoms. Such alkyl groups in
particular include methyl ("Me"), ethyl ("Et"), n-propyl ("Prⁿ"), *iso*-propyl ("Prⁱ"),
n-butyl ("Buⁿ"), *sec*-butyl ("Bu^s"), *tert*-butyl ("Bu^t"), pentyl and hexyl. Where
appropriate, such alkyl groups may be substituted by one or more groups
10 selected from halo (such as fluoro, chloro, bromo), -CN, -CF₃, -OH, -OCF₃,
C₂₋₆ alkenyl, C₃₋₆ alkynyl, C₁₋₆ alkoxy, aryl and di-C₁₋₆ alkylamino.

As used herein, the term "alkoxy" as a group or part of a group refers
to an alkyl ether radical, wherein the term "alkyl" is defined above. Such
15 alkoxy groups in particular include methoxy, ethoxy, n-propoxy, *iso*-propoxy,
n-butoxy, *iso*-butoxy, *sec*-butoxy and *tert*-butoxy. Where appropriate, such
alkoxy groups may be substituted by one or more groups selected from halo
(such as fluoro, chloro, bromo), -CN, -CF₃, -OH, -OCF₃, C₁₋₆ alkyl, C₂₋₆
alkenyl, C₃₋₆ alkynyl, aryl and di-C₁₋₆ alkylamino.

20

As used herein, the term "aryl" as a group or part of a group refers to a
carbocyclic aromatic radical ("Ar"). Suitably such aryl groups are 5-6
membered monocyclic groups or 8-10 membered fused bicyclic groups,
especially phenyl ("Ph"), biphenyl and naphthyl, particularly phenyl.

25

The term "halo" is used herein to describe, unless otherwise stated, a
group selected from fluorine ("fluoro"), chlorine ("chloro"), bromine ("bromo")
or iodine ("iodo").


30 The term "naphthyl" is used herein to denote, unless otherwise stated,
both naphth-1-yl and naphth-2-yl groups.

The term 'heterocyclyl' is used herein to describe, unless otherwise stated, groups comprising one or more rings which may be saturated, unsaturated or aromatic and which may independently contain one or more heteratoms in each ring. Examples of suitable heterocyclyl groups include, but are not limited to, acridinyl, azocinyl, benzimidazolyl, benzofuranyl, benzothiofuranyl, benzothiophenyl, benzoxazolyl, benzthiazolyl, benztriazolyl, benztetrazolyl, benzisoxazolyl, benzisothiazolyl, benzimidazolyl, carbazolyl, carbolinyl, chromanyl, chromenyl, cinnolyl, decahydroquinolyl, 2H, 6H-1,5,2-dithiazinyl, dihydrobenzofuranyl, furanyl, furazanyl, imidazolyl, 1H-indazolyl, indolyl, indolyl, isobenzofuranyl, isochromanyl, isoindazolyl, isoindolyl, isoquinolyl, isothiazolyl, isoxazolyl, naphthyridinyl, oxadiazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, oxazolyl, pyrimidinyl, phthalazinyl, pteridinyl, purinyl, pyrazinyl, pyrazolyl, pyridooxazolyl, pyridoimidazolyl, pyridothiazolyl, pyridyl, pyrrolyl, quinazolinyl, quinolyl, quinoxalinyl, tetrahydroisoquinolyl, tetrahydroquinolyl, 6H-1,2,5-thiadiazinyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, thiazolyl, thienyl, thienothiazolyl, thienooxazolyl, thienoimidazolyl, triazinyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1,2,5-triazolyl, 1,3,4-triazolyl and xanthenyl.

20


The present invention also provides, in a further aspect, a process for the preparation of a compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof, which process comprises coupling a compound of formula (II):

25



in which R^1 , P and p are as defined in formula (I), with a compound of formula (III):

30



- 5 in which R^2 and n are as defined in formula (I) and A and B contain the appropriate functional groups which are capable of reacting together to form the urea moiety;
and optionally thereafter if appropriate:
(i) removing any protecting groups;
- 10 (ii) forming a pharmaceutically acceptable salt or solvate of the compound so formed.

Suitable examples of appropriate A and B groups include:

- 15 (a) A is $-\text{N}=\text{C}=\text{O}$ and B is NH_2 ; or
(b) A is NH_2 and B is $-\text{N}=\text{C}=\text{O}$; or
(c) A is NH_2 and B is NH_2 together with an appropriate urea forming agent.
- 20 In process (a) or (b) the reaction is carried out in an inert solvent such as dichloromethane or acetonitrile.

- In process (c) the urea forming agent can be carbonyl diimidazole or phosgene. The reaction may be performed in an inert organic solvent such as dimethylformamide, tetrahydrofuran or dichloromethane at ambient or
- 25 elevated temperature. The reaction is typically performed in the presence of a base such as triethylamine or pyridine.

- An alternative method of synthesis of the unsymmetrical urea
- 30 compounds of formula (I) is from a diaryl carbonate, *via* the corresponding carbamate. Such a methodology is described by Freer et al. (Synthetic Communications, 26(2), 331 - 349, 1996). It will be appreciated that such a

methodology could readily be adapted for the preparation of compounds of formula (I).

Those skilled in the art will appreciate that it may be necessary to protect certain groups in the synthesis of compounds of formula (I). Suitable protecting groups and methods for their attachment and removal are conventional in the art of organic chemistry, such as those described in Greene T.W. 'Protective groups in organic synthesis' New York, Wiley (1981).

Compounds of formulae (II) and (III) are commercially available or may be prepared according to known methods or analogous to known methods.

Pharmaceutically acceptable salts may be prepared conventionally by reaction with the appropriate acid or acid derivative.

Compounds of formula (I) and their pharmaceutically acceptable salts or solvates thereof have Vanilloid receptor antagonist (VR1) activity and are believed to be of potential use for the treatment or prophylaxis of certain disorders, or treatment of the pain associated with them, such as: pain, chronic pain, neuropathic pain, postoperative pain, post-rheumatoid arthritic pain, osteoarthritic pain, back pain, visceral pain, cancer pain, algnesia, neuralgia, dental pain, headache, migraine, neuropathies, carpal tunnel syndrome, diabetic neuropathy, HIV-related neuropathy, post-herpetic neuralgia, fibromyalgia, neuritis, sciatica, nerve injury, ischaemia, neurodegeneration, stroke, post stroke pain, multiple sclerosis, respiratory diseases, asthma, cough, COPD, broncho constriction, inflammatory disorders, oesophagitis, heart burn, Barrett's metaplasia, dysphagia, gastroesophageal reflux disorder (GERD), stomach and duodenal ulcers, functional dyspepsia, irritable bowel syndrome, inflammatory bowel disease, colitis, Crohn's disease, pelvic hypersensitivity, pelvic pain, menstrual pain, renal colic, urinary incontinence, cystitis, burns, itch, psoriasis, pruritis, emesis (hereinafter referred to as the "Disorders of the Invention").

Thus the invention also provides a compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof, for use as a therapeutic substance, in particular in the treatment or prophylaxis of the Disorders of the
5 Invention.

In particular, the invention provides a compound of formula (I) or a pharmaceutically acceptable salt thereof or a solvate thereof for use in the treatment or prophylaxis of pain and urinary incontinence.

10

The invention further provides a method of treatment or prophylaxis of the Disorders of the Invention, in mammals including humans, which comprises administering to the sufferer a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt or solvate
15 thereof.

20

The present invention also provides a pharmaceutical composition, which comprises a compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof and a pharmaceutically acceptable carrier.

A pharmaceutical composition of the invention, which may be prepared by admixture, suitably at ambient temperature and atmospheric pressure, is usually adapted for oral, parenteral, rectal administration or intravesical administration to the bladder and, as such, may be in the form of
25 tablets, capsules, oral liquid preparations, powders, granules, lozenges, reconstitutable powders, injectable or infusable solutions, suspensions or suppositories. Orally administrable compositions are generally preferred.

Tablets and capsules for oral administration may be in unit dose form,
30 and may contain conventional excipients, such as binding agents, fillers, tableting lubricants, disintegrants and acceptable wetting agents. The tablets may be coated according to methods well known in normal pharmaceutical practice.

Oral liquid preparations may be in the form of, for example, aqueous or oily suspension, solutions, emulsions, syrups or elixirs, or may be in the form of a dry product for reconstitution with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives such as suspending agents, emulsifying agents, non-aqueous vehicles (which may include edible oils), preservatives, and, if desired, conventional flavourings or colourants.

For parenteral administration, fluid unit dosage forms are prepared utilising a compound of the invention or pharmaceutically acceptable salt or solvate thereof and a sterile vehicle. The compound, depending on the vehicle and concentration used, can be either suspended or dissolved in the vehicle. In preparing solutions, the compound can be dissolved for injection and filter sterilised before filling into a suitable vial or ampoule and sealing. Advantageously, adjuvants such as a local anaesthetic, preservatives and buffering agents are dissolved in the vehicle. To enhance the stability, the composition can be frozen after filling into the vial and the water removed under vacuum. Parenteral suspensions are prepared in substantially the same manner, except that the compound is suspended in the vehicle instead of being dissolved, and sterilization cannot be accomplished by filtration. The compound can be sterilised by exposure to ethylene oxide before suspension in a sterile vehicle. Advantageously, a surfactant or wetting agent is included in the composition to facilitate uniform distribution of the compound.

The composition may contain from 0.1% to 99% by weight, preferably from 10 to 60% by weight, of the active material, depending on the method of administration.

The dose of the compound used in the treatment of the aforementioned disorders will vary in the usual way with the seriousness of the disorders, the weight of the sufferer, and other similar factors. For systemic administration, dosage levels from 0.01mg to 100mg per

kilogramme of body weight are useful in the treatment of pain. However, as a general guide suitable unit doses may be 0.05 to 1000 mg, more suitably 0.05 to 20, 20 to 250, or 0.1 to 500.0 mg, for example 0.2 to 5 and 0.1 to 250 mg; and such unit doses may be administered more than once a day, for example
5 two or three a day, so that the total daily dosage is in the range of about 0.5 to 1000 mg; and such therapy may extend for a number of weeks or months.

When administered in accordance with the invention, no unacceptable toxicological effects are expected with the compounds of the
10 invention.

The following Examples illustrate the preparation of the compounds of the invention and are not intended to be limiting in any way.

15 **Abbreviations:**

AIBN – azoisobutyronitrile

DMF – dimethylformamide

MgSO₄ – magnesium sulfate

THF - tetrahydrofuran

20

Description 1

1-(2-Aminoethyl)indoline (D1)

A mixture of indoline (15g, 0.126mol) and 2-bromoethylamine
25 hydrobromide (12.9g, 0.063mol) in toluene was heated at reflux for 18h. After cooling solvent was removed under reduced pressure and the residue dissolved in water. Basification using aqueous potassium carbonate was followed by solvent extraction using dichloromethane. Organic phase was separated, dried over MgSO₄, filtered and concentrated under reduced
30 pressure to leave an oil. Chromatography on silica gel eluting with dichloromethane and methanol (gradient elution, maximum 20%) afforded the title compound as yellow oil (5.45g, 27%).

Description 2**4-Methylindoline (D2)**

To a solution of 4-methylindole (1g, 7.6mmol) in glacial acetic acid
5 (10ml) was added portionwise sodium cyanoborohydride (1.44g, 0.023mol)
over 15mins under an argon atmosphere. Stirring was continued for 18h and
water (100ml) added. Basification using aqueous sodium hydroxide was
followed by solvent extraction using dichloromethane. Organic phase was
separated, dried (MgSO₄), filtered and concentrated under reduced pressure
10 to leave an oil (0.98g, 97%).

Description 3**1-(2-Aminoethyl)-4-methylindoline (D3)**

15 The title compound was prepared from 4-methylindoline (D2) using
the procedure outlined for Description 1 (0.63g, 96%).

Description 4**5-Fluoroindoline (D4)**

20

The title compound was prepared from 5-fluoroindole using the
procedure outlined for Description 2 (4.1g, 82%).

Description 525 **1-(2-Aminoethyl)-5-fluoroindoline (D5)**

The title compound was prepared from 5-fluoroindoline (D4) using the
procedure outlined for Description 1 (2.5g, 92%).

30

Description 6**5-Methylindoline (D6)**

The title compound was prepared from 5-methylindole using the
5 procedure outlined for Description 2 (2.5g, 50%).

Description 7**1-(2-Aminoethyl)-5-methylindoline (D7)**

10 The title compound was prepared from 5-methylindoline (D6) using
the procedure outlined for Description 1 (1.35g, 82%).

Description 8**4-Fluoroindoline (D8)**

15

The title compound was prepared from 4-fluoroindole using the
procedure outlined for Description 2 (1.24g, 61%).

Description 9

20 **1-(2-Aminoethyl)-4-fluoroindoline (D9)**

The title compound was prepared from 4-fluoroindoline (D8) using the
procedure outlined for Description 1 (0.7g, 97%).

25 **Description 10**

N-(2-Bromo-4-fluorophenyl)acetamide (D10)

To a solution of 2-bromo-4-fluoroaniline (7.35g, 0.039mol) and
triethylamine (11ml) in dichloromethane (30ml) was added a solution of acetyl
30 chloride (2.8ml) in dichloromethane (20ml) whilst cooling (ice bath) over a
period 20 min. Stirring was continued for 6h and then dichloromethane
partitioned with water. The dichloromethane layer was separated, dried

(MgSO₄), filtered and concentrated under reduced pressure to afford the product as a white solid (4.61g, 49%).

Description 11

5 ***N*-(2-Bromo-4-fluorophenyl)-*N*-(2-methylallyl)acetamide (D11)**

To a cooled solution (ice bath) of (D10) (4.6g, 0.021mol) in dry DMF (30ml) was added sodium hydride (60% dispersion in oil, 0.89g, 0.022mol) under an argon atmosphere. After stirring for 10min 3-bromo-2-
10 methylpropene was added and the reaction stirred at ambient temperature for 18h. DMF was removed under reduced pressure and the residue partitioned between diethyl ether and water. The organic layer was separated, dried (MgSO₄), filtered and concentrated under reduced pressure to afford an oil. Chromatography on silica gel eluting with diethyl ether afforded the product
15 as a clear oil (5.5g, 91%).

Description 12

1-(5-Fluoro-3,3-dimethyl-2,3-dihydroindol-1-yl)ethanone (D12)

20 A solution of (D11) (2.2g, 8mmol) in toluene (30ml) was treated with AIBN (catalytic amount, 30mg) followed by tributyltin hydride (2.6ml, 0.01mol) in toluene (10ml). Reaction was stirred for 20min at ambient temperature and then warmed to 50°C for 90min. After cooling, the reaction mixture was partitioned with water. The organic layer was separated, dried (MgSO₄),
25 filtered and concentrated under reduced pressure to afford an oil. Chromatography on silica gel eluting with diethyl ether and hexane (gradient elution, maximum 50%), afforded the product as a white solid (0.56g, 35%).

30

Description 13**5-Fluoro-3,3-dimethylindoline (D13)**

A solution of (D12) (0.78g, 3.8mmol) in ethanol (10ml) and 2M HCl
5 (25ml) was heated at 85⁰C for 2h and cooled. Basification using sodium
hydrogen carbonate was followed by solvent extraction using
dichloromethane. Organic phase was separated, dried (MgSO₄), filtered and
concentrated under reduced pressure to leave an oil (0.55g, 87%).

10 Description 14**2-(5-Fluoro-3,3-dimethyl-2,3-dihydroindol-1-yl)ethylamine (D14)**

The title compound was prepared from (D13) using the procedure
outlined for Description 1 (0.14g, 43%).

15

Description 15**2-(6-Methyl-3,4-dihydro-2H-quinolin-1-yl)ethylamine**

The title compound was prepared from 6-methyl-1,2,3,4-
20 tetrahydroquinoline using the procedure outlined for Description 1 (5.85g,
44%).

Description 16**(5-Methyl-2-nitrophenoxy)-acetic acid, ethyl ester (D16)**

25

A solution of 5-methyl-2-nitrophenol (10g, 0.065mol) and ethyl
bromoacetate (7.25ml) in acetone (200ml) containing powdered potassium
carbonate (9.91g) was refluxed for 18h and cooled. The solid was filtered off
and the filtrate concentrated under reduced pressure to afford the product as
30 a light yellow solid (15.22g, 98%).

Description 17**7-Methyl-4H-benzo[1,4]oxazin-3-one (D17)**

A solution of (D16) (15.2g, 0.064mol) in ethanol (80ml) and
5 cyclohexene (10ml) containing 10% palladium on charcoal (0.5g) was heated
at reflux for 27h and cooled. Catalyst was filtered off and the filtrate
concentrated under reduced pressure to afford the compound as a white solid
(6.25g, 60%).

10 Description 18**7-Methyl-3,4-dihydro-2H-benzo[1,4]oxazine (D18)**

A suspension of (D17) (6.0g, 0.037mol) in dry THF (50ml) was treated
with borane.THF complex (2.5eq., 100ml). The resulting solution was then
15 heated at reflux for 3h, cooled, basified using aqueous potassium carbonate
(10%, 150ml) and extracted using dichloromethane. Organic phase was
separated, dried (MgSO₄), filtered and concentrated under reduced pressure
to leave an oil. Chromatography on silica gel eluting with dichloromethane
afforded the product as an oil (5.4g, 98%).

20

Description 19**2-(7-Methyl-2,3-dihydrobenzo[1,4]oxazin-4-yl)ethylamine (D19)**

The title compound was prepared from 7-methyl-3,4-dihydro-2H-
25 benzo[1,4]oxazine (D18) using the procedure outlined for Description 1
(2.20g, 64%).

Description 20**(5-Fluoro-2-nitrophenoxy)-acetic acid, ethyl ester (D20)**

30

The title compound was prepared from 5-fluoro-2-nitrophenol using
the procedure outlined in Description 16 (15.17g, 98%).

Description 21**7-Fluoro-4*H*-benzo[1,4]oxazin-3-one (D21)**

The title compound was prepared from (D20) using the procedure
5 outlined for Description 17 (8.15g, 79%).

Description 22**7-Fluoro-3,4-dihydro-2*H*-benzo[1,4]oxazine (D22)**

10 The title compound was prepared from (D21) using the procedure
outlined for Description 18 (7.15g, 98%).

Description 23**2-(7-Fluoro-2,3-dihydrobenzo[1,4]oxazin-4-yl)ethylamine (D23)**

15 The title compound was prepared from (D22) using the procedure
outlined for Description 19 (2.15g, 56%).

Example 1

20 **N-(2-Bromophenyl)-*N'*-[2-(7-fluoro-2,3-dihydrobenzo[1,4]oxazin-4-yl)ethyl]urea (E1)**

To a solution of 2-(7-fluoro-2,3-dihydrobenzo[1,4]oxazin-4-yl)ethylamine (D23) (0.1g, 0.5mmol) in dichloromethane (3ml) was added 2-bromophenyl isocyanate (101mg, 0.5mmol) in dichloromethane (2ml).
25 Reaction was stirred for 18h and the precipitated solid was filtered off and dried to afford the title compound as a white solid (190mg, 96%).

¹H NMR (400MHz, CDCl₃) δ7.96 (d, 1H), 7.51 d, 1H), 7.28 (m, 1H), 6.94 (m, 1H), 6.67 (m, 2H), 6.54 (m, 2H), 4.98 (br, 1H), 4.21 (m, 2H), 3.50 (m, 2H), 3.40 (m, 2H), 3.32 (m, 2H). MH⁺ 394, 396.
30

The following examples (Table 1) were prepared using a similar procedure to that outlined for E1 with the appropriate amine and isocyanate.

5

10

15

20

25

30

Table 1

Example	MH⁺
<i>N</i>-(4-Fluorophenyl)-<i>N'</i>-[2-(2,3-dihydroindol-1-yl)ethyl]urea (E2)	300
<i>N</i>-(3,4-Difluorophenyl)-<i>N'</i>-[2-(2,3-dihydroindol-1-yl)ethyl]urea (E3)	318
<i>N</i>-(2-Methyl-3-chlorophenyl)-<i>N'</i>-[2-(2,3-dihydroindol-1-yl)ethyl]urea (E4)	330, 332
<i>N</i>-(2,3-Dichlorophenyl)-<i>N'</i>-[2-(2,3-dihydroindol-1-yl)ethyl]urea (E5)	314, 316
<i>N</i>-(2-Bromophenyl)-<i>N'</i>-[2-(2,3-dihydroindol-1-yl)ethyl]urea (E6)	360, 362
<i>N</i>-(2-(Trifluoromethoxy)phenyl)-<i>N'</i>-[2-(2,3-dihydroindol-1-yl)ethyl]urea (E7)	366
<i>N</i>-(2-Iodophenyl)-<i>N'</i>-[2-(2,3-dihydroindol-1-yl)ethyl]urea (E8)	408
<i>N</i>-Phenyl-<i>N'</i>-[2-(4-methyl-2,3-dihydroindol-1-yl)ethyl]urea (E9)	296
<i>N</i>-(4-Fluorophenyl)-<i>N'</i>-[2-(4-methyl-2,3-dihydroindol-1-yl)ethyl]urea (E10)	314
<i>N</i>-(2-Chlorophenyl)-<i>N'</i>-[2-(4-methyl-2,3-dihydroindol-1-yl)ethyl]urea (E11)	330, 332
<i>N</i>-(3,4-Difluorophenyl)-<i>N'</i>-[2-(4-methyl-2,3-dihydroindol-1-yl)ethyl]urea (E12)	332
<i>N</i>-(3-Acetylphenyl)-<i>N'</i>-[2-(4-methyl-2,3-dihydroindol-1-yl)ethyl]urea (E13)	338
<i>N</i>-(2-Methyl-3-chlorophenyl)-<i>N'</i>-[2-(4-methyl-2,3-dihydroindol-1-yl)ethyl]urea (E14)	344, 346

<i>N</i> -(2,3-Dichlorophenyl)- <i>N'</i> -[2-(4-methyl-2,3-dihydroindol-1-yl)ethyl]urea (E15)	364, 366
<i>N</i> -(2,5-Dichlorophenyl)- <i>N'</i> -[2-(4-methyl-2,3-dihydroindol-1-yl)ethyl]urea (E16)	364, 366
<i>N</i> -(2-Bromophenyl)- <i>N'</i> -[2-(4-methyl-2,3-dihydroindol-1-yl)ethyl]urea (E17)	374, 376
<i>N</i> -(2-(Trifluoromethoxy)phenyl)- <i>N'</i> -[2-(4-methyl-2,3-dihydroindol-1-yl)ethyl]urea (E18)	380
<i>N</i> -(2-Iodophenyl)- <i>N'</i> -[2-(4-methyl-2,3-dihydroindol-1-yl)ethyl]urea (E19)	422
<i>N</i> -(2,5-Dichlorophenyl)- <i>N'</i> -[2-(5-methyl-2,3-dihydroindol-1-yl)ethyl]urea (E20)	364, 366
<i>N</i> -(2-Bromophenyl)- <i>N'</i> -[2-(5-methyl-2,3-dihydroindol-1-yl)ethyl]urea (E21)	374, 376
<i>N</i> -(4-Fluorophenyl)- <i>N'</i> -[2-(5-methyl-2,3-dihydroindol-1-yl)ethyl]urea (E22)	314
<i>N</i> -Phenyl- <i>N'</i> -[2-(4-fluoro-2,3-dihydroindol-1-yl)ethyl]urea (E23)	300
<i>N</i> -(4-Fluorophenyl)- <i>N'</i> -[2-(4-fluoro-2,3-dihydroindol-1-yl)ethyl]urea (E24)	318
<i>N</i> -(2-Chlorophenyl)- <i>N'</i> -[2-(4-fluoro-2,3-dihydroindol-1-yl)ethyl]urea (E25)	334, 336
<i>N</i> -(3,4-Difluorophenyl)- <i>N'</i> -[2-(4-fluoro-2,3-dihydroindol-1-yl)ethyl]urea (E26)	336
<i>N</i> -(3-Acetylphenyl)- <i>N'</i> -[2-(4-fluoro-2,3-dihydroindol-1-yl)ethyl]urea (E27)	342
<i>N</i> -(2-Methyl-3-chlorophenyl)- <i>N'</i> -[2-(4-fluoro-2,3-dihydroindol-1-yl)ethyl]urea (E28)	348, 350
<i>N</i> -(1-Naphthyl)- <i>N'</i> -[2-(4-fluoro-2,3-dihydroindol-1-yl)ethyl]urea (E29)	350

<i>N</i> -(2,3-Dichlorophenyl)- <i>N'</i> -[2-(4-fluoro-2,3-dihydroindol-1-yl)ethyl]urea (E30)	368, 370
<i>N</i> -(2,5-Dichlorophenyl)- <i>N'</i> -[2-(4-fluoro-2,3-dihydroindol-1-yl)ethyl]urea (E31)	368, 370
<i>N</i> -(2-Bromophenyl)- <i>N'</i> -[2-(4-fluoro-2,3-dihydroindol-1-yl)ethyl]urea (E32)	378, 380
<i>N</i> -(2-(Trifluoromethoxy)phenyl)- <i>N'</i> -[2-(4-fluoro-2,3-dihydroindol-1-yl)ethyl]urea (E33)	383
<i>N</i> -Phenyl- <i>N'</i> -[2-(5-fluoro-2,3-dihydroindol-1-yl)ethyl]urea (E34)	300
<i>N</i> -(4-Fluorophenyl)- <i>N'</i> -[2-(5-fluoro-2,3-dihydroindol-1-yl)ethyl]urea (E35)	318
<i>N</i> -(2-Chlorophenyl)- <i>N'</i> -[2-(5-fluoro-2,3-dihydroindol-1-yl)ethyl]urea (E36)	334, 336
<i>N</i> -(3,4-Difluorophenyl)- <i>N'</i> -[2-(5-fluoro-2,3-dihydroindol-1-yl)ethyl]urea (E37)	336
<i>N</i> -(3-Acetylphenyl)- <i>N'</i> -[2-(5-fluoro-2,3-dihydroindol-1-yl)ethyl]urea (E38)	342
<i>N</i> -(2-Methyl-3-chlorophenyl)- <i>N'</i> -[2-(5-fluoro-2,3-dihydroindol-1-yl)ethyl]urea (E39)	334, 336
<i>N</i> -(1-Naphthyl)- <i>N'</i> -[2-(5-fluoro-2,3-dihydroindol-1-yl)ethyl]urea (E40)	350
<i>N</i> -(2,5-Dichlorophenylphenyl)- <i>N'</i> -[2-(5-fluoro-2,3-dihydroindol-1-yl)ethyl]urea (E41)	368, 370
<i>N</i> -(2-Bromophenyl)- <i>N'</i> -[2-(5-fluoro-2,3-dihydroindol-1-yl)ethyl]urea (E42)	378, 380
<i>N</i> -(2-(Trifluoromethoxy)phenyl)- <i>N'</i> -[2-(5-fluoro-2,3-dihydroindol-1-yl)ethyl]urea (E43)	384
<i>N</i> -(2-Iodophenyl)- <i>N'</i> -[2-(5-fluoro-2,3-dihydroindol-1-yl)ethyl]urea (E44)	426

<i>N</i> -(2-Bromophenyl)- <i>N'</i> -[2-(6-methyl-3,4-dihydro-2 <i>H</i> -quinolin-1-yl)ethyl]urea (E45)	388, 390
<i>N</i> -(4-Fluorophenyl)- <i>N'</i> -[2-(6-methyl-3,4-dihydro-2 <i>H</i> -quinolin-1-yl)ethyl]urea (E46)	329
<i>N</i> -(2-Methyl-3-chlorophenyl)- <i>N'</i> -[2-(6-methyl-3,4-dihydro-2 <i>H</i> -quinolin-1-yl)ethyl]urea (E47)	358, 360
<i>N</i> -(1-Naphthyl)- <i>N'</i> -[2-(6-methyl-3,4-dihydro-2 <i>H</i> -quinolin-1-yl)ethyl]urea (E48)	360
<i>N</i> -(2,3-Dichlorophenyl)- <i>N'</i> -[2-(6-methyl-3,4-dihydro-2 <i>H</i> -quinolin-1-yl)ethyl]urea (E49)	378, 380
<i>N</i> -(2,5-Dichlorophenyl)- <i>N'</i> -[2-(6-methyl-3,4-dihydro-2 <i>H</i> -quinolin-1-yl)ethyl]urea (E50)	378, 380
<i>N</i> -(4-Fluorophenyl)- <i>N'</i> -[2-(7-methyl-2,3-dihydrobenzo[1,4]oxazin-4-yl)ethyl]urea (E51)	330
<i>N</i> -(2-Bromophenyl)- <i>N'</i> -[2-(7-methyl-2,3-dihydrobenzo[1,4]oxazin-4-yl)ethyl]urea (E52)	390, 392
<i>N</i> -(4-Fluorophenyl)- <i>N'</i> -[2-(7-fluoro-2,3-dihydrobenzo[1,4]oxazin-4-yl)ethyl]urea (E53)	334
<i>N</i> -(2-Bromophenyl)- <i>N'</i> -[2-(5-fluoro-3,3-dimethyl-2,3-dihydroindol-1-yl)ethyl]urea (E54)	406, 408
<i>N</i> -(1-Naphthyl)- <i>N'</i> -[2-(5-fluoro-3,3-dimethyl-2,3-dihydroindol-1-yl)ethyl]urea (E55)	378
<i>N</i> -(3-Methylcinnolin-5-yl)- <i>N'</i> -[2-(7-fluoro-2,3-dihydrobenzo[1,4]oxazin-4-yl)ethyl]urea (E56)	382
<i>N</i> -(1-Methylisoquinolin-5-yl)- <i>N'</i> -[2-(7-fluoro-2,3-dihydrobenzo[1,4]oxazin-4-yl)ethyl]urea (E57)	381
<i>N</i> -(3-Methylisoquinolin-5-yl)- <i>N'</i> -[2-(7-fluoro-2,3-dihydrobenzo[1,4]oxazin-4-yl)ethyl]urea (E58)	381

Pharmacological Data

In vitro assay

As referenced above, the compounds of the invention are vanilloid
5 receptor (VR1) antagonists and hence have useful pharmaceutical properties.
Vanilloid receptor (VR1) antagonist activity can be confirmed and
demonstrated for any particular compound by use of conventional methods,
for example those disclosed in standard reference texts such as D. Le Bars,
M. Gozarin and S. W. Cadden, Pharmacological Reviews, 2001, 53(4), 597-
10 652] or such other texts mentioned herein.


The screen used for the compounds of this invention was based upon
a FLIPR based calcium assay, similar to that described by Smart et al. (British
Journal of Pharmacology, 2000, 129, 227-230). Transfected astrocytoma
1321N1 cells, stably expressing human VR1, were seeded into FLIPR plates
15 at 25,000cells/well (96-well plate) and cultured overnight.

The cells were subsequently loaded in medium containing 4 μ M Fluo-3
AM (Molecular Probes) for 2 hours, at room temperature, in the dark. The
plates were then washed 4 times with Tyrode containing 1.5mM calcium,
without probenecid. The cells were pre-incubated with compound or buffer
20 control at room temperature for 30 minutes. Capsaicin (Sigma) was then
added to the cells. Compounds having antagonist activity against the human
VR1 were identified by detecting differences in fluorescence when measured
after capsaicin addition, compared with no compound buffer controls. Thus,
for example, in the buffer control capsaicin addition results in an increase in
25 intracellular calcium concentration resulting in fluorescence. A compound
having antagonist activity blocks the capsaicin binding to the receptor, there is
no signalling and therefore no increase in intracellular calcium levels and
consequently lower fluorescence. pK_b values are generated from the IC₅₀
values using the Cheng-Prusoff equation.

30 All compounds tested by the above methodology had pK_b > 6,
preferred compounds [Examples 1, 3, 10, 11, 12, 14-17, 19, 20-25, 27-33, 37,
39-47, 49, 50, 52, 54, 56, 57 and 58] had a pK_b > 7.0.

Claims

1. A compound of formula (I) or a pharmaceutically acceptable salt or
 5 solvate thereof:




- 10 wherein:

P is phenyl, naphthyl or heterocyclyl;

R^1 is selected from -H, halo, alkyl, alkoxy, cycloalkyl, aralkyl, aralkoxy, cycloalkylalkyl, cycloalkylalkoxy, -CN, -NO₂, -OH, -OCF₃, -CF₃, -NR⁵R⁶, -S(O)_mR⁷, -S(O)₂NR⁵R⁶, -OS(O)₂R⁷, -OS(O)₂CF₃, -O(CH₂)_xNR⁵R⁶, -

- 15 C(O)CF₃, -C(O)alkyl, -C(O)cycloalkyl, -C(O)aralkyl, -C(O)Ar, -C(O)(CH₂)_xOR⁷, -C(O)(CH₂)_xNR⁵R⁶, -C(O)alkoxy, -C(O)NR⁵R⁶, -(CH₂)_xC(O)alkoxy, -(CH₂)_xOC(O)R⁷, -(CH₂)_xOR⁷, -(CH₂)_xR⁵R⁶, -(CH₂)_xC(O)NR⁵R⁶, -(CH₂)_xN(R⁵)C(O)R⁷, -(CH₂)_xS(O)₂NR⁵R⁶, -(CH₂)_xN(R⁵)S(O)₂R⁷, -ZAr, -(CH₂)_xS(O)₂R⁷, -(OCH₂)_xS(O)₂R⁷, -
 20 N(R⁵)S(O)₂R⁷, -N(R⁵)C(O)R⁷, -(CH₂)_xN(R⁵)S(O)₂R⁷, -(CH₂)_xN(R⁵)C(O)R⁷ or -(CH₂)_xC(O)alkyl;

R^2 is the group:




- 25 X is a bond, C, O or NR⁸;

- R^3 is selected from $-H$, halo, alkyl, alkoxy, cycloalkyl, aryl, aralkyl, aralkoxy, cycloalkylalkyl, cycloalkylalkoxy, $-CN$, $-NO_2$, $-OH$, $-OCF_3$, $-CF_3$, $-NR^5R^6$, $-S(O)_mR^7$, $-S(O)_2NR^5R^6$, $-OS(O)_2R^7$, $-OS(O)_2CF_3$, $-O(CH_2)_xNR^5R^6$, $-C(O)CF_3$, $-C(O)alkyl$, $-C(O)cycloalkyl$, $-C(O)aralkyl$, $-C(O)Ar$, $-C(O)(CH_2)_xOR^7$, $-C(O)(CH_2)_xNR^5R^6$, $-C(O)alkoxy$, $-C(O)NR^5R^6$, $-(CH_2)_xC(O)alkoxy$, $-(CH_2)_xOC(O)R^7$, $-(CH_2)_xOR^7$, $-(CH_2)_xR^5R^6$, $-(CH_2)_xC(O)NR^5R^6$, $-(CH_2)_xN(R^5)C(O)R^7$, $-(CH_2)_xS(O)_2NR^5R^6$, $-(CH_2)_xN(R^5)S(O)_2R^7$, $-ZAr$, $-(CH_2)_xS(O)_2R^7$, $-(OCH_2)_xS(O)_2R^7$, $-N(R^5)S(O)_2R^7$, $-N(R^5)C(O)R^7$, $-(CH_2)_xN(R^5)S(O)_2R^7$, $-(CH_2)_xN(R^5)C(O)R^7$
- or $-(CH_2)_xC(O)alkyl$;
- R^4 is hydrogen or alkyl;
- R^5 and R^6 may be the same or different and represent H or alkyl or R^5 and R^6 together with the atoms to which they are attached form a C_3 - ϵ azacycloalkane, $C_{3-6}(2-oxo)azacycloalkane$ ring or C_{5-8} polymethylene chain
- optionally interrupted by heteroatoms such as O or $-NR^8$;
- Z is O, S or NR^8 ;
- R^7 is alkyl or aryl;
- R^8 is hydrogen, alkyl or aryl;
- n is 2, 3, 4, 5 or 6;
- p is 0, 1, 2, 3 or 4;
- q is 0, 1, 2 or 3;
- r is 0, 1 or 2; and
- x is 0, 1, 2, 3, 4, 5 or 6.

2. A compound of formula (I) as claimed in claim 1, in which P is phenyl.
3. A compound of formula (I) as claimed in claim 1 or claim 2, in which n is 2.

30

4. A compound of formula (I) as claimed in any preceding claim, in which R^2 is



5


5. A compound of formula (I) as claimed in claim 4, in which R^2 is dihydroindolyl, tetrahydroquinolyl or dihydrobenzo[1,4]oxazinyl.

- 10 6. A compound according to any of the preceding claims in which R^3 is halo or alkyl.

7. A compound according to claim 1 which is compound Example number E1 to E58 or a pharmaceutically acceptable salt or solvate thereof.


15

8. A process for the preparation of a compound of formula (I) or a pharmaceutically acceptable salt or solvate thereof, which process comprises coupling a compound of formula (II):



20

in which R^1 , P and p are as defined in formula (I) with a compound of formula (III):



25

in which R^2 and n are as defined in formula (I) and A and B contain the appropriate functional groups which are capable of reacting together to form the urea moiety;

and optionally thereafter if appropriate:

- 5 (i) removing any protecting groups;
(ii) forming a pharmaceutically acceptable salt or solvate of the compound so formed.

9. A compound of formula (I), as claimed in any one of claims 1 to 7 for
10 use in therapy.

10. A pharmaceutical composition which comprises a compound of
formula (I) as claimed in any one of claims 1 to 7 and a pharmaceutically
acceptable carrier or excipient.

15

11. The use of a compound of formula (I) as claimed in claim 1, or a
pharmaceutically acceptable salt or solvate thereof, in the manufacture of a
medicament for the treatment or prophylaxis of disorders in which an
antagonist of the vanilloid receptor (VR1) is beneficial.

20

12. A method of treating disorders or diseases in which an antagonist
of the vanilloid receptor (VR1) is beneficial which comprises administering a
safe and therapeutically effective amount to a patient in need thereof of a
compound of formula (I) as claimed in claim 1, or a pharmaceutically effective
25 salt or solvate thereof.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
3 July 2003 (03.07.2003)

PCT

(10) International Publication Number
WO 2003/053945 A3

(51) International Patent Classification⁷: **C07D 265/36**,
A61K 31/395, A61P 43/00, C07D 209/14, 215/12, 413/12

(74) Agent: **GIDDINGS, Peter, John**; GlaxoSmithKline,
Corporate Intellectual Property CN925.1, 980 Great West
Road, Brentford, Middlesex TW8 9GS (GB).

(21) International Application Number:
PCT/GB2002/005812

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC,
VN, YU, ZA, ZM, ZW.

(22) International Filing Date:
19 December 2002 (19.12.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0130550.7 20 December 2001 (20.12.2001) GB

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK,
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*):
SMITHKLINE BEECHAM P.L.C. [GB/GB]; 980
Great West Road, Brentford, Middlesex TW8 9GS (GB).

Published:

— with international search report


(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **RAMI, Harshad**,
Kantlilal [GB/GB]; GlaxoSmithKline, New Frontiers Sci-
ence Park South, Third Avenue, Harlow, Essex CM19 5AW
(GB). **THOMPSON, Mervyn** [GB/GB]; GlaxoSmithK-
line, New Frontiers Science Park South, Third Avenue,
Harlow, Essex CM19 5AW (GB).

(88) Date of publication of the international search report:
11 March 2004

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: UREA DERIVATIVES AND THEIR USE AS VANILLOID RECEPTOR ANTAGONISTS



(57) Abstract: Compounds of formula (I) or a pharmaceutically acceptable salt or solvate thereof: (I) wherein, R¹, R², P, p and n are as defined in the specification, a process for preparing such compounds, a pharmaceutical composition containing such compounds and the use of such compounds in medicine.

WO 2003/053945 A3

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB 02/05812

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D265/36 A61K31/395 A61P43/00 C07D209/14 C07D215/12
C07D413/12

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHEMICAL ABSTRACTS, vol. 112, no. 2, 8 January 1990 (1990-01-08) Columbus, Ohio, US; abstract no. 138852s, HOGALE, M.B. ET AL: "Preparation of 5-methyl-N-propylamino-2,3-dihydroindole derivatives." XP002248859 abstract & J. INDIAN CHEM. SOC., vol. 66, no. 7, - 1989 pages 484-486, -& DATABASE CAPLUS 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; DN 112:138852, XP002248860 compound with RN 125178-82-7 --- -/--	1

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

24 July 2003

Date of mailing of the international search report

28/08/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Van Bijlen, H

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB 02/05812

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 3 247 211 A (MAX A. WEAVER ET AL.) 19 April 1966 (1966-04-19) * example 21(a) * ---	1
A	US 5 840 720 A (ING-JUN CHEN) 24 November 1998 (1998-11-24) column 3 -column 4 ---	1,10,11
A	WO 98 20867 A (THE UNITED STATES OF AMERICA) 22 May 1998 (1998-05-22) page 11 -----	1,10,11

INTERNATIONAL SEARCH REPORT

International application No.
PCT/GB 02/05812

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claim 12 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 02/05812

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 3247211	A	19-04-1966	BE 661212 A DE 1292280 B FR 1387387 A GB 1062272 A	16-07-1965 10-04-1969 19-05-1965 22-03-1967
US 5840720	A	24-11-1998	NONE	
WO 9820867	A	22-05-1998	AU 5460598 A WO 9820867 A1	03-06-1998 22-05-1998